arXiv:1207.0198v1 [math.NT] 1 Jul 2012

SEMI-ORDINARY p-STABILIZATION OF THE SIEGEL EISENSTEIN SERIES OF ARBITRARY GENUS

HISA-AKI KAWAMURA

1. Introduction

For each even integer $\kappa > 2$, as a typical example of modular form with respect to the elliptic modular group $\mathrm{SL}_2(\mathbb{Z})$, the (holomorphic) Eisenstein series is defined by

$$E_\kappa(z)(=E_\kappa^{(1)}(z)) := \frac{\zeta(1-\kappa)}{2} \sum_{\substack{(c,d)\in\mathbb{Z}^2,\\ \gcd(c,d)=1,\, d>0}} (cz+d)^{-\kappa},$$

where $z \in \mathfrak{H}_1 = \{z \in \mathbb{C} \,|\, \mathrm{Im}\, z > 0\}$ and $\zeta(s)$ is Riemann's zeta function. It is well-known that E_κ is a holomorphic modular form of weight κ with respect to $\mathrm{SL}_2(\mathbb{Z})$, which possesses the Fourier expansion

$$E_\kappa(z) = \frac{\zeta(1-\kappa)}{2} + \sum_{m=1}^{\infty} \sigma_{\kappa-1}(m) q^m,$$

where $\sigma_{\kappa-1}(m) = \sum_{0<d|m} d^{\kappa-1}$ and $q = \exp(2\pi\sqrt{-1}z)$. In addition, E_κ forms a non-cuspidal Hecke eigenform such that Hecke's L-function $L(s,\, E_\kappa)$ is taken of the form

$$L(s,\, E_\kappa) = \zeta(s)\zeta(s-\kappa+1).$$

Let $p \neq 2$ be a rational prime. Put

$$E_\kappa^*(z) := E_\kappa(z) - p^{\kappa-1}E_\kappa(pz). \tag{1}$$

Then we easily see that E_κ^* is also a non-cuspidal Hecke eigenform of weight κ with respect to the congruence subgroup $\Gamma_0(p) \subset \mathrm{SL}_2(\mathbb{Z})$ of level p, and the Fourier expansion is taken of the form

$$E_\kappa^*(z) = \frac{\zeta^{(p)}(1-\kappa)}{2} + \sum_{m=1}^{\infty} \sigma_{\kappa-1}^{(p)}(m)\, q^m, \tag{2}$$

Date: May 30th, 2012 (Preliminary Version).

2010 *Mathematics Subject Classification.* 11F46 (Primary); 11F30, 11F33 (Secondary).

Key words and phrases. Siegel modular forms, Eisenstein series, p-adic analytic family.

where $\zeta^{(p)}(s) := (1-p^{-s})\zeta(s)$ and $\sigma^{(p)}_{\kappa-1}(m) := \sum_{\substack{0<d|m,\\ \gcd(d,p)=1}} d^{\kappa-1}$. It turns out that for E^*_κ, all the Hecke eigenvalues outside p agree with those of E_κ, but the eigenvalue of Atkin's U_p-operator (cf. [AL70, Mi06]) is $\sigma^{(p)}_{\kappa-1}(p) = 1$. Namely, we have

$$L(s,\, E^*_\kappa) = \zeta(s)\zeta^{(p)}(s-\kappa+1).$$

This type of p-adic normalization selecting half the Euler factor at p is often said to be the *ordinary p-stabilization.* Here we should mention that every Fourier coefficient of E^*_κ depends on the weight κ p-adically analytically. Indeed, for each positive integer m, Fermat's little theorem implies that the function $\kappa \mapsto \sigma^{(p)}_{\kappa-1}(m)$ can be extended to an analytic function defined on the ring of p-adic integers $\mathbb{Z}_p$. In addition, the constant term $\zeta^{(p)}(1-\kappa)/2$ is also interpolated by the p-adic zeta function in the sense of Kubota-Leopoldt and Iwasawa p-adically. Thus, we obtain the following:

Fact 1.1 (cf. [Hi93] §7)**.** Let $\Lambda := \mathbb{Z}_p[[X]]$ be the one-variable power series ring over $\mathbb{Z}_p$, and $\mathcal{L} = \mathrm{Frac}(\Lambda)$ the field of fractions of Λ, respectively. Let $\omega : \mathbb{Z}_p^\times \twoheadrightarrow \mu_{p-1}(\mathbb{Q}_p)$ be the Teichmüller character, where $\mu_{p-1}(\mathbb{Q}_p) \simeq (\mathbb{Z}/p\mathbb{Z})^\times$ is the torsion subgroup of $\mathbb{Z}_p^\times$, regarded as a Dirichlet character of conductor p. Then for each integer $0 \le a < p-1$, there exists a formal Fourier expansion

$$\mathbf{E}_{\omega^a}(X)(= \mathbf{E}^{(1)}_{\omega^a}(X)) = \sum_{m=0}^{\infty} \mathbf{a}_m(\omega^a;\, X)q^m \in \mathcal{L}[[q]]$$

such that for each even integer $\kappa > 2$ with $\kappa \equiv a \pmod{p-1}$, we have

$$\mathbf{E}_{\omega^a}((1+p)^\kappa - 1) = E^*_\kappa.$$

Therefore, for a given positive even integer $\kappa_0 > 2$, an infinite collection of ordinary p-stabilized Eisenstein series' $\{E^*_\kappa\}_\kappa$, in which the weights κ vary under the inequality $\kappa \ge \kappa_0$ and the congruence relation

$$\kappa \equiv \kappa_0 \pmod{(p-1)p^n}$$

for some positive integer n sufficiently large, is a p-adic analytic family of Hecke eigenforms of level p.

The aim of the present article is to know how to formulate these phenomena in the case of Siegel modular forms, that is, automorphic form on the symplectic group $\mathrm{Sp}_{2g}(\mathbb{Q})$ of arbitrary genus $g \ge 1$. To be more precise, for the Siegel Eisenstein series $E^{(g)}_\kappa$ with respect to $\mathrm{Sp}_{2g}(\mathbb{Z})$ (to be defined in §2 below), we define a certain p-stabilization by means of the action of a linear combination of $U^i_{p,g}$ $(i = 0, 1, \cdots, g)$, where $U_{p,g}$ denotes the generalized U_p-operator (cf. Equation (10) below), so that the eigenvalue of $U_{p,g}$ is equal to 1. Throughout the article, we refer to it as a *semi-ordinary* p-stabilization according to the same tradition as Skinner-Urban [SU06] (see also [Pa00] and [K10]). Moreover, we derive an explicit form of the resulting Fourier expansion which can be viewed as a natural generalization of the equation

(2) (cf. Theorem 3.2 below). As a consequence of the above issues, we show in Theorems 3.5 and 3.6 below that semi-ordinary p-stabilized Siegel Eisenstein series' taken above form a p-adic analytic family of Siegel modular forms of genus g and level p in which the weights vary p-adically.

Acknowledgements. This research was partially supported by the JSPS Grant-in-Aid for Scientific Research with the assigned ID 23224001 and 22340001.

Notation. We denote by $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$, and $\mathbb{C}$ the ring of integers, fields of rational numbers, real numbers and complex numbers, respectively. Let $\overline{\mathbb{Q}}$ denote the algebraic closure of $\mathbb{Q}$ sitting inside $\mathbb{C}$. We put $e(x) = \exp(2\pi\sqrt{-1}x)$ for $x \in \mathbb{C}$. For a rational prime p, we denote by $\mathbb{Q}_p$, $\mathbb{Z}_p$ and $\mathbb{Z}_p^\times$ the field of p-adic numbers, the ring of integers and the group of units, respectively. Hereinafter, we fix an algebraic closure $\overline{\mathbb{Q}}_p$ of $\mathbb{Q}_p$ and an embedding of $\overline{\mathbb{Q}}$ into $\overline{\mathbb{Q}}_p$ once for all. Let v_p denote the p-adic valuation on $\overline{\mathbb{Q}}_p$ normalized so that $v_p(p) = 1$, and e_p the continuous additive character of $\overline{\mathbb{Q}}_p$ such that $e_p(x) = e(x)$ for all $x \in \mathbb{Q}$. Let $\mathbb{C}_p$ be the completion of $\overline{\mathbb{Q}}_p$ with respect to v_p. We also fix, once for all, an isomorphism $\mathbb{C} \xrightarrow{\sim} \mathbb{C}_p$ such that the diagram

$$\begin{tikzcd}
\mathbb{C} \arrow[r, "\sim"] & \mathbb{C}_p \\
\overline{\mathbb{Q}} \arrow[u, hook] \arrow[r, hook] & \overline{\mathbb{Q}}_p \arrow[u, hook]
\end{tikzcd}$$

is commutative.

For each integer $g \geq 1$, let GSp_{2g} be the group of symplectic similitudes defined over $\mathbb{Q}$, that is,

$$\mathrm{GSp}_{2g} := \left\{ M \in \mathrm{GL}_{2g} \mid {}^tMJ_{2g}M = \mu(M)J_{2g} \text{ for some } \mu(M) \in \mathbb{G}_m \right\},$$

where $J_{2g} = \left(\begin{smallmatrix} 0_g & 1_g \\ -1_g & 0_g \end{smallmatrix}\right)$ with the $g \times g$ unit (resp. zero) matrix 1_g (resp. 0_g), and Sp_{2g} the derived group of GSp_{2g} characterized by the exact sequence

$$1 \to \mathrm{Sp}_{2g} \to \mathrm{GSp}_{2g} \xrightarrow{\mu} \mathbb{G}_m \to 1,$$

respectively. We note that every real point $M = \left(\begin{smallmatrix} A & B \\ C & D \end{smallmatrix}\right) \in \mathrm{GSp}_{2g}(\mathbb{R})$ with $\mu(M) > 0$, where A, B, C, $D \in \mathrm{Mat}_{g\times g}(\mathbb{R})$, acts on the Siegel upper half-space

$$\mathfrak{H}_g := \left\{ Z = X + \sqrt{-1}\,Y \in \mathrm{Mat}_{g\times g}(\mathbb{C}) \,\middle|\, {}^tZ = Z,\, Y > 0\,(\text{positive-definite}) \right\}$$

of genus g via the linear transformation $Z \mapsto M\langle Z\rangle = (AZ+B)(CZ+D)^{-1}$. If F is a function on $\mathfrak{H}_g$, then for each $\kappa \in \mathbb{Z}$, we define a natural action of M on F by

$$(F|_\kappa M)(Z) := \mu(M)^{g\kappa - g(g+1)/2} \det(CZ+D)^{-\kappa} F(M\langle Z\rangle).$$

For each integer $N \geq 1$, put

$$\Gamma_0(N)^{(g)} := \left\{ M \in \mathrm{Sp}_{2g}(\mathbb{Z}) \;\middle|\; M \equiv \begin{pmatrix} * & * \\ 0_g & * \end{pmatrix} \pmod N \right\},$$
$$\Gamma_1(N)^{(g)} := \left\{ M \in \mathrm{Sp}_{2g}(\mathbb{Z}) \;\middle|\; M \equiv \begin{pmatrix} * & * \\ 0_g & 1_g \end{pmatrix} \pmod N \right\}.$$

For each $\kappa \in \mathbb{Z}$, let us denote by $\mathscr{M}_\kappa(\Gamma_1(N))^{(g)}$ the space of *classical Siegel modular forms* of genus g, weight κ and level N, that is, $\mathbb{C}$-valued holomorphic functions F on $\mathfrak{H}_g$ satisfying the following conditions:

(i) $F|_\kappa M = F$ for any $M \in \Gamma_1(N)^{(g)}$.

(ii) For each $M \in \mathrm{Sp}_{2g}(\mathbb{Z})$, the function $F|_\kappa M$ possesses a Fourier expansion of the form

$$(F|_\kappa M)(Z) = \sum_{T \in \mathrm{Sym}_g^*(\mathbb{Z})} A_T(F|_\kappa M)\, e(\mathrm{tr}(TZ)),$$

where let $\mathrm{Sym}_g^*(\mathbb{Z})$ denote the set of all half-integral symmetric matrices of degree g over $\mathbb{Z}$, namely,

$$\mathrm{Sym}_g^*(\mathbb{Z}) := \{T = (t_{ij}) \in \mathrm{Sym}_g(\mathbb{Q}) \,|\, t_{ii},\, 2t_{ij} \in \mathbb{Z}\ (1 \le i < j \le g)\},$$

and $\mathrm{tr}(*)$ the trace, respectively. Then it is satisfied that

$$A_T(F|_\kappa M) = 0 \text{ unless } T \geq 0 \text{ (positive-semidefinite)}$$

for all $M \in \mathrm{Sp}_{2g}(\mathbb{Z})$.

A modular form $F \in \mathscr{M}_\kappa(\Gamma_1(N))^{(g)}$ is said to be *cuspidal* (or a *cusp form*) if it is satisfied that $A_T(F|_\kappa M) = 0$ unless $T > 0$ for all $M \in \mathrm{Sp}_{2g}(\mathbb{Z})$. We denote by $\mathscr{S}_\kappa(\Gamma_1(N))^{(g)}$ the subspace of $\mathscr{M}_\kappa(\Gamma_1(N))^{(g)}$ consisting of all cuspidal forms. For each Dirichlet character χ modulo N, we denote by $\mathscr{M}_\kappa(\Gamma_0(N),\, \chi)^{(g)}$ (resp. $\mathscr{S}_\kappa(\Gamma_0(N),\, \chi)^{(g)}$) the subspace of $\mathscr{M}_\kappa(\Gamma_1(N))^{(g)}$ (resp. $\mathscr{S}_\kappa(\Gamma_1(N))^{(g)}$) consisting of all forms F with Nebentypus character χ, that is,

$$F|_\kappa M = \chi(\det D) F \quad \text{for any } M = \left(\begin{smallmatrix} A & B \\ C & D \end{smallmatrix}\right) \in \Gamma_0(N)^{(g)}.$$

In particular, if χ is the principal character, we naturally write

$$\mathscr{M}_\kappa(\Gamma_0(N))^{(g)} = \mathscr{M}_\kappa(\Gamma_0(N), \chi)^{(g)} \text{ and } \mathscr{S}_\kappa(\Gamma_0(N))^{(g)} = \mathscr{S}_\kappa(\Gamma_0(N),\, \chi)^{(g)},$$

respectively.

For each $T = (t_{ij}) \in \mathrm{Sym}_g^*(\mathbb{Z})$ and $Z = (z_{ij}) \in \mathfrak{H}_g$, put

$$\mathbf{q}^T := e(\mathrm{tr}(TZ)) = \prod_{i=1}^{g} q_{ii}^{t_{ii}} \prod_{i<j\le g} q_{ij}^{2t_{ij}},$$

where $q_{ij} = e(z_{ij})\, (1 \le i \le j \le n)$. It is easy to see that for each $F \in \mathscr{M}_\kappa(\Gamma_1(N))^{(g)}$, the usual Fourier expansion

$$F(Z) = \sum_{\substack{T \in \mathrm{Sym}_g^*(\mathbb{Z}), \\ T \geq 0}} A_T(F)\, \mathbf{q}^T$$

belongs to the ring $\mathbb{C}[\, q_{ij}^{\pm 1} \,|\, 1 \le i < j \le g \,][[q_{11}, \cdots, q_{gg}]]$. For each commutative ring R, we write

$$R[[\mathbf{q}]]^{(g)} := R[\, q_{ij}^{\pm 1} \,|\, 1 \le i < j \le g \,][[q_{11}, \cdots, q_{gg}]],$$

in a similar fashion as the notation of the ring of formal q-expansions $R[[q]]$. In particular, if $F \in \mathscr{M}_\kappa(\Gamma_1(N))^{(g)}$ is a Hecke eigenform (i.e. a simultaneous eigenfunction of all Hecke operators whose similitude is coprime to N), then it is well-known that the field K_F obtained by adjoining all Fourier coefficients (or equivalently, all Hecke eigenvalues) of F to $\mathbb{Q}$ is a totally real number field, to which we refer as the *Hecke field* of F. Namely, we have

$$F \in K_F[[\mathbf{q}]]^{(g)} \hookrightarrow \overline{\mathbb{Q}}[[\mathbf{q}]]^{(g)} \hookrightarrow \mathbb{C}_p[[\mathbf{q}]]^{(g)}.$$

For further details on Siegel modular forms set out above, see [AZ95] or [Fr83].

2. The Siegel Eisenstein series of genus g

In this section, we review some fundamental properties of the Siegel Eisenstein series of arbitrary genus $g \ge 1$.

Let r be a positive integer. For each rational prime l, let $\mathrm{Sym}_r^*(\mathbb{Z}_l)$ denote the set of all half-integral symmetric matrices of degree r over $\mathbb{Z}_l$. For each nondegenerate $S \in \mathrm{Sym}_r^*(\mathbb{Z}_l)$, we define a formal power series $b_l(S;\, X)$ in X by

$$b_l(S;\, X) := \sum_{R \in \mathrm{Sym}_r(\mathbb{Q}_l)/\mathrm{Sym}_r(\mathbb{Z}_l)} e_l(\mathrm{tr}(SR)) X^{v_l(\mu_R)},$$

where $\mu_R = [\mathbb{Z}_l^g + \mathbb{Z}_l^g R : \mathbb{Z}_l^g]$. Put $\mathfrak{D}_S := 2^{2[r/2]} \det S$. In particular, if g is even, then $(-1)^{r/2}\mathfrak{D}_S \equiv 0,\, 1 \pmod 4$, and thus, we may decompose it into the form

$$(-1)^{r/2}\mathfrak{D}_S = \mathfrak{d}_S\, \mathfrak{f}_S^2,$$

where $\mathfrak{d}_S$ is the fundamental discriminant corresponding to the quadratic field extension $\mathbb{Q}_l(\{(-1)^{r/2}\mathfrak{D}_S\}^{1/2})/\mathbb{Q}_l$ and $\mathfrak{f}_S = \{(-1)^{r/2}\mathfrak{D}_S/\mathfrak{d}_S\}^{1/2} \in \mathbb{Z}_l$. Let $\chi_l : \mathbb{Q}_l^\times \to \{\pm 1,\, 0\}$ denote the character defined by

$$\chi_l(x) = \begin{cases} 1 & \text{if } \mathbb{Q}_l(x^{1/2}) = \mathbb{Q}_l, \\ -1 & \text{if } \mathbb{Q}_l(x^{1/2})/\mathbb{Q}_l \text{ is unramified}, \\ 0 & \text{if } \mathbb{Q}_l(x^{1/2})/\mathbb{Q}_l \text{ is ramified}. \end{cases}$$

As shown in [Ki84, Fe86, Sh94], there exists a polynomial $F_l(S;\, X) \in \mathbb{Z}[X]$ such that the formal power series $b_l(S;\, X)$ can be decomposed into

$$b_l(S;\, X) = \begin{cases} \dfrac{(1-X)\prod_{i=1}^{r/2}(1-l^{2i}X^2)}{1-\chi_l((-1)^{r/2}\det S)l^{r/2}X}\, F_l(S;\, X) & \text{if } r \text{ is even}, \\ (1-X)\prod_{i=1}^{(r-1)/2}(1-l^{2i}X^2)\, F_l(S;\, X) & \text{if } r \text{ is odd}. \end{cases} \tag{3}$$

We note that $F_l(S;\,0)=1$, and that $F_l(S;\,X)$ satisfies the functional equation

$$(4)\quad F_l(S;\,l^{-r-1}X^{-1})=\begin{cases}(l^{r+1}X^2)^{-v_l(\mathfrak{f}_S)}F_l(S;\,X) & \text{if } r \text{ is even},\\ \eta_l(S)(l^{(r+1)/2}X)^{-v_l(\mathfrak{D}_S)}F_l(S;\,X) & \text{if } r \text{ is odd},\end{cases}$$

where let $h_l(S)$ denote the Hasse invariant of S in the sense of Kitaoka, $(*,*)_l$ the Hilbert symbol defined over $\mathbb{Q}_l$, and

$$\eta_l(S):=h_l(S)(\det S,\,(-1)^{(r-1)/2}\det S)_l(-1,\,-1)_l^{(r^2-1)/8},$$

respectively (cf. Theorem 3.2 in [Ka99]). Thus, $F_l(S;\,X)$ has degree $2v_l(\mathfrak{f}_S)$ or $v_l(\mathfrak{D}_S)$ according as g is even or odd. We easily see that $F_l(uS;\,X)=F_l(S;\,X)$ for each $u\in\mathbb{Z}_l^\times$. For further details on the above issues, see [Ka99].

On the other hand, for a positive integer g, and a positive even integer κ with $\kappa>g+1$, the (*holomorphic*) *Siegel Eisenstein series* of genus g and weight κ is defined as follows: For each $Z\in\mathfrak{H}_g$, put

$$\begin{aligned}E_\kappa^{(g)}(Z) \;&:=\; 2^{-[(g+1)/2]}\zeta(1-\kappa)\prod_{i=1}^{[g/2]}\zeta(1-2\kappa+2i)\\ &\quad\times\sum_{M=\left(\begin{smallmatrix}* & *\\ C & D\end{smallmatrix}\right)\in\mathrm{P}_{2g}\cap\mathrm{Sp}_{2g}(\mathbb{Z})\backslash\mathrm{Sp}_{2g}(\mathbb{Z})}\det(CZ+D)^{-\kappa},\end{aligned}$$

where $\mathrm{P}_{2g}=\{\left(\begin{smallmatrix}* & *\\ 0_g & *\end{smallmatrix}\right)\in\mathrm{GSp}_{2g}\}$. It is well-known that $E_\kappa^{(g)}\in\mathscr{M}_\kappa(\mathrm{Sp}_{2g}(\mathbb{Z}))$ is a non-cuspidal Hecke eigenform. Indeed, let us denote the Fourier expansion by

$$E_\kappa^{(g)}(Z)=\sum_{\substack{T\in\mathrm{Sym}_g^*(\mathbb{Z}),\\ T\geq 0}}A_T(E_\kappa^{(g)})\,\mathbf{q}^T,$$

then for $T_0\in\mathrm{Sym}_g^*(\mathbb{Z})$ with $\operatorname{rank}T_0=0$, we have

$$(5)\qquad A_{T_0}(E_\kappa^{(g)})=2^{-[(g+1)/2]}\zeta(1-\kappa)\prod_{i=1}^{[g/2]}\zeta(1-2\kappa+2i).$$

More generally, if $T\in\mathrm{Sym}_g^*(\mathbb{Z})$ is taken of the form

$$T=\left(\begin{array}{c|c}T' & \\ \hline & 0_{g-r}\end{array}\right)$$

for some nondegenerate $T'\in\mathrm{Sym}_r^*(\mathbb{Z})$ with $0<r\leq g$ (i.e. $\operatorname{rank}T=r$), then we have

$$\begin{aligned}(6)\quad & A_T(E_\kappa^{(g)})\\ &=\; 2^{[(r+1)/2]-[(g+1)/2]}\prod_{i=[r/2]+1}^{[g/2]}\zeta(1-2\kappa+2i)\end{aligned}$$

$$\times\begin{cases} L\left(1-\kappa+r/2,\, \left(\frac{\mathfrak{d}_{T'}}{*}\right)\right) \displaystyle\prod_{l\mid\mathfrak{f}_{T'}} F_l(T';\, l^{\kappa-r-1}) & \text{if } r \text{ is even},\\ \displaystyle\prod_{l\mid\mathfrak{D}_{T'}} F_l(T';\, l^{\kappa-r-1}) & \text{if } r \text{ is odd},\end{cases}$$

where $L(s,\, \left(\frac{\mathfrak{d}}{*}\right))$ denotes the Dirichlet L-function associated with the Kronecker character $\left(\frac{\mathfrak{d}}{*}\right)$ of conductor $\mathfrak{d}$. For the convenience in the sequel, we make the convention that if $\operatorname{rank} T = 0$, then $\mathfrak{D}_T = \mathfrak{d}_T = \mathfrak{f}_T = 1$, and $F_l(T;\, X) = 1$. Accordingly, the equation (5) is collapsed into the preceding equation (6).

Finally in this section, we make a few remarks on the aspect of $E_\kappa^{(g)}$ as a Hecke eigenform. For each prime l, let

$$(\psi_{l,0}(E_\kappa^{(g)}), \psi_{l,1}(E_\kappa^{(g)}), \cdots, \psi_{l,g}(E_\kappa^{(g)})) \in (\mathbb{C}^\times)^{g+1}/W_g$$

denote the Satake parameter of $E_\kappa^{(g)}$ at l, where W_g is the Weyl group isomorphic to $\mathfrak{S}_g \ltimes \{\pm 1\}^g$. Then the associated spinor L-function is defined as follows:

$$L(s,\, E_\kappa^{(g)},\, \mathrm{spin}) = \prod_{l:\mathrm{prime}} \mathcal{Q}_l(E_\kappa^{(g)};\, l^{-s})^{-1},$$

where $\mathcal{Q}_l(E_\kappa^{(g)};\, Y) \in \mathbb{C}[Y]$ denotes the Hecke-Andrianov polynomial defined by

$$\begin{aligned}&\mathcal{Q}_l(E_\kappa^{(g)};\, Y)\\ &\quad := \ (1-\psi_{l,0}(E_\kappa^{(g)})Y) \prod_{r=1}^{g} \prod_{1\le i_1<\cdots<i_r\le g} (1-\psi_{l,0}(E_\kappa^{(g)})\psi_{l,i_1}(E_\kappa^{(g)})\cdots\psi_{l,i_r}(E_\kappa^{(g)})Y),\end{aligned}$$

If g is even, then for each l, we may take the Satake parameter $(\psi_{l,i}(E_\kappa^{(g)}))_{0\le i\le g}$ as

$$\psi_{l,i}(E_\kappa^{(g)}) = \begin{cases} l^{(g/2)\cdot(\kappa-g/2)-(g/2)\cdot(g/2+1)/2} & \text{if } i=0,\\ l^{-\kappa+g/2+i} & \text{if } 1\le i\le g/2,\\ l^{\kappa-g+i-1} & \text{if } g/2+1\le i\le g.\end{cases} \tag{7}$$

(cf. [Mu02]). If $g>1$ is odd, the Zharkovskaya identity (cf. [Ža74]) yields

$$L(s,\, E_\kappa^{(g)},\, \mathrm{spin}) = L(s,\, E_\kappa^{(g-1)},\, \mathrm{spin})L(s-\kappa+g,\, E_\kappa^{(g-1)},\, \mathrm{spin}),$$

and thus, for each prime l, we have

$$\psi_{l,i}(E_\kappa^{(g)}) = \begin{cases} \psi_{l,0}(E_\kappa^{(g-1)}) & \text{if } i=0,\\ \psi_{l,i}(E_\kappa^{(g-1)}) & \text{if } 1\le i\le g-1,\\ l^{\kappa-g} & \text{if } i=g.\end{cases} \tag{8}$$

Therefore, regardless of the parity of g, $L(s,\, E_\kappa^{(g)},\, \mathrm{spin})$ is expressed as a product of $\zeta(s)$'s.

3. Main results

Let us fix a rational prime $p \neq 2$ once for all. In this section, for each positive integer g, we introduce a certain p-stabilization of the Siegel Eisenstein series $E_\kappa^{(g)} \in \mathscr{M}_\kappa(\mathrm{Sp}_{2g}(\mathbb{Z}))$, which can be viewed as a natural generalization of the ordinary p-stabilization of $E_\kappa^{(1)}$ (cf. (1)).

To begin with, we introduce the following two polynomials in X and Y:

$$\begin{aligned} \mathcal{P}_{p,g}(X,\,Y) &:= (1-p^gXY)\prod_{i=1}^{[g/2]}(1-p^{2g-2i+1}X^2Y), \\ \mathcal{R}_{p,g}(X,\,Y) &:= \prod_{j=1}^{g}(1-p^{j(2g-j+1)/2}X^jY). \end{aligned}$$

In addition, let us denote by $\widetilde{\mathcal{R}}_{p,g}(X,\,Y)$ the reflected polynomial of $\mathcal{R}_{p,g}(X,\,Y)$ with respect to Y, that is,

$$\begin{aligned} \widetilde{\mathcal{R}}_{p,g}(X,\,Y) &:= Y^g\,\mathcal{R}_{p,g}(X,\,Y^{-1}) \\ &:= \prod_{j=1}^{g}(Y-p^{j(2g-j+1)/2}X^j). \end{aligned} \tag{9}$$

Example 3.1. When $g=1$ and 2, a straightforward calculation yields

$$\begin{cases} \mathcal{P}_{p,1}(X,\,Y)=\mathcal{R}_{p,1}(X,\,Y)=1-pXY, \\ \mathcal{P}_{p,2}(X,\,Y)=\mathcal{R}_{p,2}(X,\,Y)=(1-p^2XY)(1-p^3X^2Y). \end{cases}$$

If $g>2$, then $\mathcal{P}_{p,g}(X,\,Y)\neq\mathcal{R}_{p,g}(X,\,Y)$, however, $\mathcal{P}_{p,g}(X,\,1)$ divides $\mathcal{R}_{p,g}(X,\,1)$ in general.

On the other hand, recall that if $g>1$, the following Hecke operators at p are fundamental stuffs in the p-adic theory of Siegel modular forms of genus g: For each integer $N\geq 1$, put

$$U_{p,i} := \begin{cases} \Gamma_0(N)^{(g)}\,\mathrm{diag}(\underbrace{1,\cdots,1}_{i}\underbrace{p,\cdots,p}_{g-i},\underbrace{p^2,\cdots,p^2}_{i}\underbrace{p,\cdots,p}_{g-i})\,\Gamma_0(N)^{(g)}, & \text{if } 1\leq i\leq g-1, \\ \Gamma_0(N)^{(g)}\,\mathrm{diag}(\underbrace{1,\cdots,1}_{g},\underbrace{p,\cdots,p}_{g})\,\Gamma_0(N)^{(g)}, & \text{if } i=g. \end{cases}$$

We note if N is divisible by p, these operators $U_{p,1},\,\cdots,\,U_{p,g-1}$ and $U_{p,g}$ generate the dilating Hecke algebra at p acting on Siegel modular forms of level N. In particular, we are interested in the operator $U_{p,g}$ which plays a central role among them. Indeed, we easily see that if $F\in\mathscr{M}_\kappa(\Gamma_0(N))^{(g)}$ with some integers $\kappa\geq 0$ and $N\geq 1$, then

$$F|_\kappa U_{p,g} \in \begin{cases} \mathscr{M}_\kappa(\Gamma_0(N))^{(g)}, & \text{if } p\,|\,N, \\ \mathscr{M}_\kappa(\Gamma_0(Np))^{(g)}, & \text{if } p\nmid N, \end{cases}$$

and the action of $U_{p,g}$ commutes with those of all Hecke operators at each prime $l \neq p$. In addition, if $F = \sum_{T \geq 0} A_T(F)\mathbf{q}^T \in \mathbb{C}_p[[\mathbf{q}]]^{(g)}$, then the action of $U_{p,g}$ effects on F as

$$F|_\kappa U_{p,g} = \sum_{T \geq 0} A_{pT}(F)\mathbf{q}^T, \tag{10}$$

and thus, we may regard it as a natural generalization of Atkin's U_p-operator defined in the case where $g = 1$ (cf. [AL70, Mi06]). For further details on the operator $U_{p,g}$, see, for instance, [AZ95, Bö05].

Now, we introduce a p-stabilization of $E_\kappa^{(g)}$ as follows.

Theorem 3.2. *Let κ be a positive even integer with $\kappa > g+1$. For each prime $p \neq 2$, put*

$$(E_\kappa^{(g)})^* := \frac{\mathcal{P}_{p,g}(p^{\kappa-g-1},\, 1)}{\mathcal{R}_{p,g}(p^{\kappa-g-1},\, 1)} \cdot E_\kappa^{(g)}|_\kappa\, \widetilde{\mathcal{R}}_{p,g}(p^{\kappa-g-1},\, U_{p,g}). \tag{11}$$

Then we have

(i) *$(E_\kappa^{(g)})^* \in \mathscr{M}_\kappa(\Gamma_0(p))^{(g)}$ is a non-cuspidal Hecke eigenform such that all the eigenvalues outside p agree with those of $E_\kappa^{(g)}$. Namely, we have*

$$L^{(p)}(s,\, (E_\kappa^{(g)})^*,\, \mathrm{spin}) = L^{(p)}(s,\, E_\kappa^{(g)},\, \mathrm{spin}),$$

where $L^{(p)}(s,\, E_\kappa^{(g)},\, \mathrm{spin})$ and $L^{(p)}(s,\, (E_\kappa^{(g)})^*,\, \mathrm{spin})$ both denote the spinor L-functions with the Euler factors at p removed.

(ii) *If $0 \leq T \in \mathrm{Sym}_g^*(\mathbb{Z})$ is equivalent over $\mathbb{Z}_p$ to*

$$\left(\begin{array}{c|c} T' & \\ \hline & 0_{g-r} \end{array}\right)$$

for some nondegenerate $T' \in \mathrm{Sym}_r^*(\mathbb{Z}_p)$ with $0 \leq r \leq g$, then the T-th Fourier coefficient of $(E_\kappa^{(g)})^*$ is taken of the form

$$\begin{aligned} & A_T((E_\kappa^{(g)})^*) \\ &= 2^{[(r+1)/2]-[(g+1)/2]} \prod_{i=[r/2]+1}^{[g/2]} \zeta^{(p)}(1-2\kappa+2i) \\ &\quad \times \begin{cases} L^{(p)}(1-\kappa+r/2,\, \left(\frac{\mathfrak{d}_{T'}}{*}\right)) \displaystyle\prod_{\substack{l | \mathfrak{f}_{T'}, \\ l \neq p}} F_l(T';\, l^{\kappa-r-1}) & \textit{if } r \textit{ is even}, \\ \displaystyle\prod_{\substack{l | \mathfrak{D}_{T'}, \\ l \neq p}} F_l(T';\, l^{\kappa-r-1}) & \textit{if } r \textit{ is odd}, \end{cases} \end{aligned}$$

where $L^{(p)}(s,\, \left(\frac{\mathfrak{d}}{*}\right)) := (1 - \left(\frac{\mathfrak{d}}{p}\right) p^{-s}) L(s,\, \left(\frac{\mathfrak{d}}{*}\right))$. In particular, we have

$$(E_\kappa^{(g)})^*|_\kappa U_{p,g} = (E_\kappa^{(g)})^*.$$

Proof. The assertion (i) is obvious. The assertion (ii) has been already proved in [K10] in the case where g is even. For readers' convenience, we give a complete proof of it here, which proceeds in the same way as the one of Theorem 5.1 (and also Theorem 4.1) in [K10]. When $r = 0$, the assertion immediately follows from the equation (5) and

$$\mathcal{P}_{p,g}(p^{\kappa-g-1},\, 1) = (1 - p^{\kappa-1}) \prod_{i=1}^{[g/2]} (1 - p^{2\kappa-2i-1}).$$

Hereinafter, suppose that $r > 0$. It follows by definition that

$$\widetilde{\mathcal{R}}_{p,g}(X, Y) \;\; = \;\; \sum_{m=0}^{g} (-1)^m s_m\Big(\{p^{j(2g-j+1)/2} X^j \,|\, 1 \le j \le g\}\Big)\, Y^{g-m},$$

where for each $0 \le m \le g$, let $s_m(\{X_1, \cdots, X_g\})$ denote the m-th elementary symmetric polynomial in $X_1, \cdots, X_g$. Thus, by the equations (6), (10) and (11), we have

$$\begin{aligned} A_T((E_\kappa^{(g)})^*) &= 2^{[(r+1)/2]-[(g+1)/2]} \prod_{i=[r/2]+1}^{[g/2]} \zeta(1 - 2\kappa + 2i)\, \frac{\mathcal{P}_{p,g}(p^{\kappa-g-1},\, 1)}{\mathcal{R}_{p,g}(p^{\kappa-g-1},\, 1)} \\ &\times \sum_{m=0}^{g} (-1)^m s_m\Big(\{p^{j(2g-j+1)/2} (p^{\kappa-g-1})^j \,|\, 1 \le j \le g\}\Big)\, F_p(p^{g-m}T';\, p^{\kappa-r-1}) \\ &\times \begin{cases} L(1 - \kappa + r/2,\, \left(\frac{\mathfrak{d}_{T'}}{*}\right)) \displaystyle\prod_{\substack{l \,|\mathfrak{f}_{T'}, \\ l \ne p}} F_l(T';\, l^{\kappa-r-1}) & \text{if } r \text{ is even,} \\ \displaystyle\prod_{\substack{l \,|\mathfrak{D}_{T'}, \\ l \ne p}} F_l(T';\, l^{\kappa-r-1}) & \text{if } r \text{ is odd.} \end{cases} \end{aligned}$$

Thus, to prove the assertion (ii), it suffices to show that if $p \ne 2$, then for each nondegenerate $T' \in \mathrm{Sym}_r^*(\mathbb{Z}_p)$ with $0 < r \le g$, the following equation holds valid:

$$\begin{aligned} &\sum_{m=0}^{r} (-1)^m s_m\Big(\{p^{j(2r-j+1)/2} X^j \,|\, 1 \le j \le r\}\Big)\, F_p(p^{r-m}T';\, X) \\ &\qquad = \begin{cases} \big(1 - \chi_p((-1)^{r/2} \det T')\, p^{r/2} X\big)\, \dfrac{\mathcal{R}_{p,r}(X,\, 1)}{\mathcal{P}_{p,r}(X,\, 1)} & \text{if } r \text{ is even,} \\ \dfrac{\mathcal{R}_{p,r}(X,\, 1)}{\mathcal{P}_{p,r}(X,\, 1)} & \text{if } r \text{ is odd.} \end{cases} \end{aligned} \tag{12}$$

Indeed, the desired assertion follows from the equation (12) immediately since its right-hand side is invariant under $T' \mapsto pT'$ regardless of the parity of r. On the other hand, Theorem 1 in [Ki86] implies that for each nondegenerate $T' \in \mathrm{Sym}_r^*(\mathbb{Z}_p)$, the equation

$$\sum_{m=0}^{\infty} F_p(p^m T';\, X) Y^m = \frac{\mathcal{S}_p(T';\, X,\, Y)}{(1-Y)\mathcal{R}_{p,r}(X,\, Y)}$$

holds for some polynomial $\mathcal{S}_p(T';\, X,\, Y) \in \mathbb{Z}[X,\, Y]$ (see also Theorem 6 in [BS87]). Since the preceding equation yields

$$\mathcal{S}_p(T';\, X,\, 1) = \sum_{m=0}^{r} (-1)^m s_m\Big(\{p^{j(2r-j+1)/2} X^j \mid 1 \le j \le r\}\Big)\, F_p(p^{r-m} T';\, X),$$

we may reformulate the equation (12) as

(13)

$$\mathcal{S}_p(T';\, X,\, 1) = \begin{cases} \big(1 - \chi_p((-1)^{r/2} \det T')\, p^{r/2} X\big)\, \dfrac{\mathcal{R}_{p,r}(X,\, 1)}{\mathcal{P}_{p,r}(X,\, 1)} & \text{if } r \text{ is even,} \\[2ex] \dfrac{\mathcal{R}_{p,r}(X,\, 1)}{\mathcal{P}_{p,r}(X,\, 1)} & \text{if } r \text{ is odd.} \end{cases}$$

When $r = 1$, we easily see that $F_p(t;\, X) = \sum_{i=0}^{v_p(t)} (pX)^i$ for each nonzero $t \in \mathbb{Z}_p$. Thus, we have

$$F_p(pt;\, X) - pX F_p(t;\, X) = 1 = \frac{\mathcal{R}_{p,1}(X, 1)}{\mathcal{P}_{p,1}(X, 1)}.$$

When $r > 1$, for a given T, let $\mathfrak{i}(T)$ denote the least integer m such that $p^m T^{-1} \in \mathrm{Sym}_r^*(\mathbb{Z}_p)$. It turns out that for each nondegenerate $T \in \mathrm{Sym}_2^*(\mathbb{Z}_p)$, the polynomial $F_p(T;\, X)$ admits the explicit form

$$\begin{aligned} & F_p(T;\, X) \\ & \quad = \sum_{i=0}^{\mathfrak{i}(T)} (p^2 X)^i \left\{ \sum_{j=0}^{v_p(\mathfrak{f}_T)-i} (p^3 X^2)^j - \chi_p(-\det T)\, pX \sum_{j=0}^{v_p(\mathfrak{f}_T)-i-1} (p^3 X^2)^j \right\} \end{aligned}$$

(cf. [Kau59, Ka99]). Thus a simple calculation yields that

$$F_p(p^2 T;\, X) - (p^2 X + p^3 X^2) F_p(pT;\, X) + p^5 X^3 F_p(T;\, X) = 1 - \chi_p(-\det T) pX,$$

and hence, the equation (13) also holds for $r = 2$. Now, let us assume that $r \ge 3$. We note that every nondegenerate $T \in \mathrm{Sym}_r^*(\mathbb{Z}_p)$ is equivalent to the form

$$\left(\begin{array}{cc|c} t_1 & & \\ & t_2 & \\ \hline & & T_3 \end{array}\right)$$

over $\mathbb{Z}_p$, where $t_1,\, t_2 \in \mathbb{Z}_p - \{0\}$ and $T_3 \in \mathrm{Sym}_{r-2}^*(\mathbb{Z}_p) \cap \mathrm{GL}_{r-2}(\mathbb{Q}_p)$ admit $v_p(t_1) \ge v_p(t_2) \ge \mathfrak{i}(T_3)$. It follows from Theorem 4.1 in [Ka99] that

$$\frac{\mathcal{S}_p(T;\, X,\, 1)}{1 - \chi_p((-1)^{r/2} \det T) p^{r/2} X} \quad = \quad \frac{\mathcal{S}_p(T_3;\, p^2 X,\, 1)}{1 - \chi_p((-1)^{r/2-1} \det T_3) p^{r/2+1} X}$$

$$\times \frac{(1-p^{(r-1)(r+2)/2}X^{r-1})(1-p^{r(r+1)/2}X^{r})}{1-p^{r+1}X^2}$$

if r is even, and

$$\mathcal{S}_p(T;\, X,\, 1) = \mathcal{S}_p(T_3;\, p^2X,\, 1) \cdot \frac{(1-p^{(r-1)(r+2)/2}X^{r-1})(1-p^{r(r+1)/2}X^{r})}{1-p^{r+2}X^2}$$

if r is odd. (See also §3 of [Ka01].) Thus, by induction on r, we have proved that the desired equation (13) holds in general. □

Remark 3.3. The statement (i) of Theorem 3.2 insists that the assignment $E_\kappa^{(g)} \mapsto (E_\kappa^{(g)})^*$ can be regarded as a p-stabilization generalizing the ordinary p-stabilization of $E_\kappa^{(1)}$ (cf. §1). When $g = 2$, Skinner-Urban [SU06] dealt with a similar type of p-stabilization so that the resulting eigenvalue of $U_{p,2}$ is a p-adic unit. Accordingly, we refer to $(E_\kappa^{(g)})^*$ as a *semi-ordinary* p-stabilization of $E_\kappa^{(g)}$ even in the case $g > 2$. However, it should be emphasized that $(E_\kappa^{(g)})^*$ does not admit the ordinary condition at p in the sense of Hida (cf. [Hi02, Hi04]). Indeed, by equations (7) and (8), there is no way to produce from $E_\kappa^{(g)}$ to a simultaneous eigenfunction of $U_{p,1} \cdots,\, U_{p,g-1},\, U_{p,g}$ such that all the eigenvalues are p-adic units. It is because the fact that the spherical representation of $\mathrm{GSp}_{2g}(\mathbb{Q}_p)$ arising from $E_\kappa^{(g)}$ is a degenerate principal series.

Remark 3.4. Let $\mathcal{Q}_p(E_\kappa^{(g)};\, Y) \in \mathbb{C}[Y]$ be the Hecke-Andrianov polynomial of $E_\kappa^{(g)}$ at p (cf. §2), and put

$$\mathcal{Q}_p^*(E_\kappa^{(g)};\, Y) := \frac{\mathcal{Q}_p(E_\kappa^{(g)};\, Y)}{(1-\psi_{p,0}(E_\kappa^{(g)})Y)(1-Y)}.$$

We easily see that $\mathcal{R}_{p,g}(p^{\kappa-g-1},\, Y)$ divides $\mathcal{Q}_p^*(E_\kappa^{(g)};\, Y)$. Thus, we may deduce from Theorem 3.2 (ii) that

$$(E_\kappa^{(g)})^* = \frac{\mathcal{P}_{p,g}(p^{\kappa-g-1},\, 1)}{\mathcal{Q}_p^*(E_\kappa^{(g)};\, 1)} \cdot E_\kappa^{(g)}|_\kappa\, \widetilde{\mathcal{Q}}_p^*(E_\kappa^{(g)};\, U_{p,g}), \tag{14}$$

where $\widetilde{\mathcal{Q}}_p^*(E_\kappa^{(g)};\, Y) := Y^{2^{g-1}}\, \mathcal{Q}_p^*(E_\kappa^{(g)};\, Y^{-1})$. We note that the right-hand side of the preceding equation looks quite similar to the p-stabilization in the sense of Courtieu-Panchishkin, which also admits the semi-ordinarity at p (cf. Remark 2.5 in [CP04]). Indeed, these two p-stabilizations are the same up to constant multiples.

As a straightforward conclusion of Theorem 3.2 above, we have

Theorem 3.5. *For each positive integer g, and each integer $0 \leq a < p-1$, there exists a formal Fourier expansion*

$$\mathbf{E}^{(g)}_{\omega^a}(X) = \sum_{\substack{T \in \mathrm{Sym}^*_g(\mathbb{Z}),\\ T \geq 0}} \mathbf{A}_T(\omega^a;\, X)\, \mathbf{q}^T \in \mathcal{L}[[\mathbf{q}]]^{(g)},$$

where ω is the Teichmüller character and $\mathcal{L} = \mathrm{Frac}(\Lambda)$, such that for each even integer $\kappa > g+1$ with $\kappa \equiv a \pmod{p-1}$, we have

$$\mathbf{E}^{(g)}_{\omega^a}((1+p)^\kappa - 1) = (E^{(g)}_\kappa)^*.$$

Proof. For each $x \in \mathbb{Z}_p^\times$, put $\langle x \rangle = \omega(x)^{-1}x \in 1 + p\mathbb{Z}_p$. Namely, together with the Teichmüller character ω, this projection induces the product decomposition $\mathbb{Z}_p^\times \xrightarrow{\sim} \mu_{p-1}(\mathbb{Q}_p) \times 1 + p\mathbb{Z}_p$, $x \mapsto (\omega(x),\, \langle x \rangle)$. In addition, for each $x \in 1 + p\mathbb{Z}_p$, put $s(x) := \log_p(x)/\log_p(1+p)$, where $\log_p$ is the p-adic logarithm function, and thus we have $s : 1 + p\mathbb{Z}_p \xrightarrow{\sim} \mathbb{Z}_p$. On the other hand, as is well-known from the work of Kubota-Leopoldt and Iwasawa (cf. [Hi93] §3.4-5), for a given quadratic Dirichlet character ψ and an integer $0 \leq a < p-1$, there exists $\mathbf{\Phi}(\psi\omega^a;\, X) \in \mathcal{L}$ such that

$$\mathbf{\Phi}(\psi\omega^a;\, (1+p)^\kappa - 1) = L^{(p)}(1-\kappa,\, \psi\omega^{a-\kappa})$$

for each integer $\kappa > 1$. Then for each $T = \begin{pmatrix} T' & \\ & 0_{g-r} \end{pmatrix} \in \mathrm{Sym}^*_g(\mathbb{Z})$ with $T' \in \mathrm{Sym}^*_r(\mathbb{Z}) \cap \mathrm{GL}_r(\mathbb{Q})$ and $0 \leq r \leq g$, we define $\mathbf{A}_T(\omega^a;\, X) \in \mathcal{L}$ as follows:

$$\begin{aligned}
&\mathbf{A}_T(\omega^a;\, X)\\
&\quad = \ 2^{[(r+1)/2]-[(g+1)/2]} \prod_{i=[r/2]+1}^{[g/2]} \mathbf{\Phi}(\omega^{2a-2i};\, (1+p)^{-2i}(1+X)^2 - 1)\\
&\qquad \times \begin{cases} \mathbf{\Phi}(\left(\frac{\mathfrak{d}_{T'}}{*}\right) \omega^{a-r/2};\, (1+p)^{-r/2}(1+X) - 1) & \\ \qquad \times \displaystyle\prod_{\substack{l \mid \mathfrak{f}_{T'},\\ l \neq p}} F_l(T';\, \omega^a(l) l^{-r-1}(1+X)^{s(\langle l \rangle)}) & \text{if } r \text{ is even,}\\ \displaystyle\prod_{\substack{l \mid \mathfrak{D}_{T'},\\ l \neq p}} F_l(T';\, \omega^a(l) l^{-r-1}(1+X)^{s(\langle l \rangle)}) & \text{if } r \text{ is odd.} \end{cases}
\end{aligned}$$

By Theorem 3.2 (ii), we easily see that for each even integer $\kappa > g+1$ with $\kappa \equiv a \pmod{p-1}$, we have

$$\mathbf{A}_T(\omega^a;\, (1+p)^\kappa - 1) = A_T((E^{(g)}_\kappa)^*).$$

This completes the proof. $\square$

By Berthelot's construction (cf. [deJ96] §7), we may associate Λ to a $\mathbb{C}_p$-rigid analytic variety $\mathcal{X}_\Lambda$, whose $\mathbb{C}_p$-valued points are given by

$$\mathcal{X}_\Lambda(\mathbb{C}_p) = \mathrm{Hom}_{cont}(\mathcal{A}(\mathcal{X}_\Lambda), \mathbb{C}_p),$$

where $\mathcal{A}(\mathcal{X}_\Lambda)$ is the topological ring consisting of all $\mathbb{C}_p$-rigid analytic functions on $\mathcal{X}_\Lambda$. We note that a natural ring homomorphism $\Lambda \to \mathcal{A}(\mathcal{X}_\Lambda)$ induces an isomorphism

$$\mathcal{X}_\Lambda(\mathbb{C}_p) \overset{\sim}{\to} \mathrm{Hom}_{cont}(\Lambda, \mathbb{C}_p).$$

For each positive integer κ and each character $\varepsilon : 1 + p\mathbb{Z}_p \to \overline{\mathbb{Q}}_p^\times$ having p-power order, the evaluation map from Λ to $\mathbb{C}_p$, $X \mapsto \varepsilon(1+p)(1+p)^\kappa - 1$ defines a point $P_{\kappa,\varepsilon} \in \mathcal{X}_\Lambda(\mathbb{C}_p)$ which is said to be *arithmetic*. In particular, if ε is trivial (i.e. ε has exact order 1), we will suppress it from the notation and write simply P_κ instead of $P_{\kappa,\varepsilon}$. Accordingly, every positive integer κ is identified with an arithmetic point $P_\kappa \in \mathcal{X}_\Lambda(\mathbb{C}_p)$. Therefore, Theorem 3.5 above is reformulated as follows:

Theorem 3.6. *Let g be a positive integer. For each positive even integer $\kappa_0 > g + 1$, let $U \subseteq \mathcal{X}_\Lambda(\mathbb{C}_p)$ be an affinoid neighborhood containing P_{κ_0}. Then there exists a formal Fourier expansion $\mathbf{E}_U^{(g)} \in \mathcal{A}(U)[[\mathbf{q}]]^{(g)}$ such that for each $P_\kappa \in U$ with $\kappa \geq \kappa_0$, we have*

$$\mathbf{E}_U^{(g)}(P_\kappa) = (E_\kappa^{(g)})^*.$$

Namely, $\{(E_\kappa^{(g)})^*\}_{P_\kappa \in U}$ is a semi-ordinary p-adic analytic family of non-cuspidal Siegel modular forms (more precisely, Hecke eigenforms) of genus g and level p.

Remark 3.7. Panchishkin [Pa00] has already constructed a similar p-adic family of Siegel Eisenstein series' of arbitrary genus $g \geq 2$, however, its siblings are twisted by a Dirichlet character whose conductor is divisible by p. In that case, we may derive the p-adic interpolation properties of Fourier coefficients only for $0 < T \in \mathrm{Sym}_g^*(\mathbb{Z})$ with $\gcd(p, \det(T)) = 1$. On the contrary, by virtue of the description of the semi-ordinary p-stabilization for $E_\kappa^{(g)}$ (cf. Theorem 3.2), we resolve the p-adic interpolation problem for the whole Fourier expansion with no restriction.

When $g = 2$, Takemori [Ta12] independently proved that for each integer $r \geq 1$ and each prime-to-p integer $N \geq 1$, a similar statement remains valid for the Siegel Eisenstein series of genus 2 and level Np^r with a primitive Nebentypus character. In the present article, for simplicity, we have just dealt with the case where $N = 1$, but $g \geq 1$ is any given integer. We note that by taking into account the presence of a non-trivial Nebentypus character from the beginning, the method we use is extendable to the Siegel Eisenstein series of genus g and level $N > 1$ as well.

References

[AZ95] A.N. Andrianov and V.G. Zhuravlëv, *Modular forms and Hecke operators*, Transl. of Math. Monogr., 145. AMS, Providence, RI, 1995.

[AL70] A.O.L. Atkin and J. Lehner, *Hecke operators on* $\Gamma_0(m)$, Math. Ann. **185** (1970), 134–160.

[Bö05] S. Böcherer, *On the Hecke operator U(p). With an appendix by Ralf Schmidt*, J. Math. Kyoto Univ. **45** (2005), no.4, 807–829.

[BS87] S. Böcherer and F. Sato, *Rationality of certain formal power series related to local densities*, Comment. Math. Univ. St.Paul. **36** (1987), no.1, 53–86.

[CP04] M. Courtieu and A. Panchishkin, *Non-Archimedean L-Functions and Arithmetical Siegel Modular Forms*, Lecture Notes in Math. vol.1471, Springer-Verlag, Berlin, 2004.

[deJ96] A.J. de Jong, *Crystalline Dieudonné module theory via formal and rigid geometry*, Inst. Hautes Études Sci. Publ. Math. **82** (1996), 5–96; Inst. Hautes Études Sci. Publ. Math. **87** (1998), 175.

[Fe86] P. Feit, *Poles and residues of Eisenstein series for symplectic and unitary groups*, Mem. Amer. Math. Soc. **61** (1986), no.346, iv+89 pp.

[Fr83] E. Freitag, *Siegelsche Modulfunktionen*, Grundlehren Math. Wiss., vol.254, Springer-Verlag, Berlin, 1983.

[Hi93] H. Hida, *Elementary theory of L-functions and Eisenstein series*, London Math. Soc. Student Text **26**, Cambridge University Press, Cambridge, 1993.

[Hi02] H. Hida, *Control theorems for coherent sheaves on Shimura varieties of PEL-type*, J. Inst. Math. Jussieu **1** (2002), 1–76.

[Hi04] H. Hida, *p-adic automorphic forms on Shimura varieties*, Springer Monographs in Math., Springer-Verlag, New York, 2004.

[Ik01] T. Ikeda, *On the lifting of elliptic modular forms to Siegel cusp forms of degree* $2n$, Ann. of Math. **154** (2001), no.3, 641–681.

[Ka99] H. Katsurada, *An explicit formula for Siegel series*, Amer. J. Math. **121** (1999), 415–452.

[Ka01] H. Katsurada, *Euler factor of a certain Dirichlet series attached to Siegel Eisenstein series*, Abh. Math. Sem. Univ. Hamburg **71** (2001), 81–90.

[Kau59] G. Kaufhold, *Dirichletsche Reihe mit Funktionalgleichung in der Theorie der Modulfunktion 2. Grades*, Math. Ann. **137** (1959), 454–476.

[K10] H. Kawamura, *On certain constructions of p-adic families of Siegel modular forms of even genus*, `http://arxiv.org/abs/1011.6476`.

[Ki84] Y. Kitaoka, *Dirichlet series in the theory of quadratic forms*, Nagoya Math. J. **92** (1984), 73–84.

[Ki86] Y. Kitaoka, *Local densities of quadratic forms and Fourier coefficients of Eisenstein series*, Nagoya Math. J. **103** (1986), 149–160.

[Mi06] T. Miyake, *Modular forms*, Springer Monogr. in Math., Springer-Verlag, Berlin, 2006.

[Mu02] K. Murakawa, *Relations between symmetric power L-functions and spinor L-functions attached to Ikeda lifts*, Kodai Math. J. **25** (2002), 61–71.

[Pa00] A. Panchishkin, *On the Siegel-Eisenstein measure and its applications*, Israel J. Math. **120** (2000), Part B, 467–509.

[Sh94] G. Shimura, *Euler products and Fourier coefficients of automorphic forms on symplectic groups*, Invent. Math. **116** (1994), no.1–3, 531–576.

[SU06] C. Skinner and E. Urban, *Sur les déformations p-adiques de certaines représentations automorphes*, J. Inst. Math. Jussieu **5** (2006), no.4, 626–698.

[Ta12] S. Takemori, *p-adic Siegel-Eisenstein series of degree two*, J. Number Theory **132** (2012), no.6, 1203–1264.

[Tay88] R.L. Taylor, *Congruences between modular forms*, Ph.D thesis, Princeton University, 1988.

[Ža74] N.A. Žarkovskaja, *The Siegel operator and Hecke operators*, Funkcional. Anal. i Priložen. **8** (1974), no.2, 30–38.

Department of Mathematics, Hokkaido University, Kita 10 Nishi 8, Kita-Ku, Sapporo, 060-0810 Hokkaido, Japan
E-mail address: `hkawamura@sci.hokudai.ac.jp`